

Adult Vaccines

-Influenza (Flu)

- Recommended 1 dose annually for all ages between the months of October and November
- Ages 65 and older receive a higher dose of the vaccine

-Pneumococcal 13 (pneumonia)

- Recommended for patients age 65 and older and high risk
- Should be administered 6 to 12 months before Pneumococcal 23

-Pneumococcal 23 (pneumonia)

- Recommended for patients age 65 and older and high risk
- Should be administered 6 to 12 months after Pneumococcal 13

-Tdap & Td (Tetanus, Diphtheria, Pertussis (whooping cough))

- One time dose of Tdap then boost with Td every 10 years

-Shingrex (Shingles)

- Recommended for patients age 50 and older
- 2 dose series administered 2-6 months apart

-HPV (Cervical cancer)

- Recommended ages 11 to 26
- 2 to 3 dose series

****Vaccine Information Sheets (VIS) from the CDC are available in our office, online, and will be given to the patient before each vaccine is given. These have additional information about the vaccine the patient will be receiving.**

**** If you have any questions about vaccination, please talk to your health care professional and visit these websites below.**

Resources

CDC: www.cdc.gov/vaccines/adults

CDC: www.cdc.gov/vaccinesafety

American Academy of Family Physicians: www.aafp.org

National Institutes of Health: www.nih.gov

American College of Physicians: www.acponline.org

Children & Adolescent Vaccines

-Hep B (Hepatitis B)

Dose 1: Birth (before hospital discharge)

Dose 2: 1 to 2 months old Dose 3: 6 to 18 months old

-RV (Rotavirus)

Dose 1: 2 months old

Dose 2: 4 months old

Dose 3: 6 months old

-DTaP (Tetanus, Diphtheria, & Pertussis (whooping cough))

Dose 1: 2 months old

Dose 2: 4 months old

Dose 3: 6 months old

Dose 4: 15-18 months old

Dose 5: 4-6 years old

-Tdap (Tetanus, Diphtheria, Pertussis(whooping cough))

Onetime dose of Tdap at 11-12 years old, boost with Td every 10 yrs

-Hib (Haemophilus influenzae type b5)

Dose 1: 2 months old

Dose 2: 4 months old

Dose 3: 6 months old

Dose 4: 12-15 months old

-Pneumococcal 13 (pneumonia)

Dose 1: 2 months old

Dose 2: 4 months old

Dose 3: 6 months old

Dose 4: 12-15 months old

-IPV (Poliovirus)

Dose 1: 2 months old

Dose 2: 4 months old

Dose 3: 6- 18 months old

Dose 4: 4-6 years old

-Influenza (Flu)

Given annually after 6 months old

First time receiving the vaccine, children 8 years and under will receive 2 doses one month apart. Every year following will be 1 dose

-MMR (Measles, Mumps, Rubella)

Dose 1: 12-15 months old

Dose 2: 4-6 years old

-Varicella (Chicken Pox)

Dose 1: 12-15 months old

Dose 2: 4-6 years old

-HepA (Hepatitis A)

2 dose series between 12 and 23 months given 6 to 18 months apart

-HPV (Cervical cancer)

Recommended ages 11 to 26; 2 to 3 dose series depending when 1st dose was received

-Meningococcal

Dose 1: 11-12 years old

Booster dose: 16-17 years old

****VFC and WyVIP Vaccines are given at our practice. VFC and WyVIP are federally funded programs that provide vaccines at no cost to children under 18 years of age within our state.**

****Vaccine Information Sheets (VIS) from the CDC are available in our office, online, and will be given to the parent/ guardian before each vaccine is given. These have additional information about the vaccine the patient will be receiving.**

****As a parent/ guardian, we understand that caring for your child is the most important thing in your life. If you have any questions about vaccination, please talk to your health care professional and visit these websites below.**

Resources:

CDC: www.cdc.gov/vaccines/parents

CDC: www.cdc.gov/vaccinesafety

American Academy of Pediatrics: www.aap.org

National Institutes of Health: www.nih.gov

American College of Physicians: www.acponline.org